arXiv:1702.06446v1 [cs.IT] 21 Feb 2017

Several Classes of Permutation Trinomials over $\mathbb{F}_{5^n}$ From Niho Exponents

Gaofei Wu [*], Nian Li [†]

Abstract

The construction of permutation trinomials over finite fields attracts people's interest recently due to their simple form and some additional properties. Motivated by some results on the construction of permutation trinomials with Niho exponents, by constructing some new fractional polynomials that permute the set of the $(q+1)$-th roots of unity in $\mathbb{F}_{q^2}$, we present several classes of permutation trinomials with Niho exponents over $\mathbb{F}_{q^2}$, where $q = 5^k$.

Index Terms Finite fields, Permutation polynomials, Trinomials, Niho exponents.

1 Introduction

Let p be a prime and n be a positive integer. Let $\mathbb{F}_q$ be a finite field of $q = p^n$ elements. We denote $\mathbb{F}_q^*$ the multiplication group of $\mathbb{F}_q$. A polynomial $f \in \mathbb{F}_q[x]$ is called a permutation polynomial if the associated polynomial mapping $f : c \mapsto f(c)$ from $\mathbb{F}_q$ to itself is a permutation of $\mathbb{F}_q$ [15]. Permutation polynomials over finite fields have important applications in cryptography [3, 18], coding theory [9], and combinatorial design theory [2]. Thus, it is important to find new permutation polynomials, both theoretically and in practice.

Permutation polynomials with few terms especially binomials and trinomials attracts people's interest recently due to their simple algebraic form and additional extraordinary properties. Recently, there are lots of constructions of permutation trinomials over finte fields, please refer to [1, 5, 6, 7, 8, 10, 11, 12, 13, 14, 19, 22].

In this paper, we focus on the construction of permutation trinomials over $\mathbb{F}_{5^n}$ from Niho exponents with the form of

$$f(x) = x + \lambda_1 x^{c_1(5^k-1)+1} + \lambda_2 x^{c_2(5^k-1)+1}, \tag{1}$$

[*]G. Wu is with the State Key Laboratory of Integrated Service Networks, Xidian University, Xi'an, 710071, China. Email: wugf@nipc.org.cn

[†]N. Li is with the Department of Informatics, University of Bergen, N-5020 Bergen, Norway. Email: nianli.2010@gmail.com.

where $n = 2k$, $1 \leq s, t \leq 5^k$, and $\lambda_1, \lambda_2 \in \{1, -1\}$. A positive integer d (always understood modulo $p^n - 1$) is a Niho exponent if $d \equiv p^j \pmod{p^{\frac{n}{2}} - 1}$ for some $j < n$ [16]. By finding some new classes of fractional polynomials that permute the set of the $(5^k + 1)$-th roots of unity in $\mathbb{F}_{5^{2k}}$, we construct several classes of permutation trinomials of the form (1) over $\mathbb{F}_{5^{2k}}$.

The remainder of this paper is organized as follows. Section 2 introduces some notations and background. In Section 3, we present some classes of permutation trinomials over $\mathbb{F}_{5^n}$ by finding some fractional polynomials that permute the set of the $(5^k + 1)$-th roots of unity in $\mathbb{F}_{5^{2k}}$. Some concluding remarks are given in Section 4.

2 Preliminaries

Let p be prime, n, k be two integers such that $k \mid n$. The trace function from $\mathbb{F}_{p^n}$ onto $\mathbb{F}_{p^k}$ is defined by

$$\mathrm{Tr}_k^n(x) = \sum_{i=0}^{n/k-1} x^{p^{ik}}, \ x \in \mathbb{F}_{p^n}.$$

Lemma 1 *Denote by* $\mathrm{Tr}(x)$ *and* $\mathrm{N}(x)$ *the trace function and norm function from* $\mathbb{F}_{q^2}$ *to* $\mathbb{F}_q$*, where* $q = 5^k$*. Then*

$$\mathrm{Tr}(x^2) = \mathrm{Tr}^2(x) - 2\mathrm{N}(x), \ \mathrm{Tr}(x^3) = \mathrm{Tr}^3(x) + 2\mathrm{N}(x)\mathrm{Tr}(x),$$

$$\mathrm{Tr}(x^4) = \mathrm{Tr}^4(x) + \mathrm{N}(x)\mathrm{Tr}^2(x) + 2\mathrm{N}^2(x),$$

$$\mathrm{Tr}(x^8) = \mathrm{Tr}^8(x) + 2\mathrm{N}(x)\mathrm{Tr}^6(x) - \mathrm{N}^3(x)\mathrm{Tr}^2(x) + 2\mathrm{N}^4(x),$$

and

$$\mathrm{Tr}(x^9) = \mathrm{Tr}^9(x) + \mathrm{N}(x)\mathrm{Tr}^7(x) + 4\mathrm{N}^4(x)\mathrm{Tr}(x) + 2\mathrm{N}^2(x)\mathrm{Tr}^5(x).$$

Lemma 1 can be proved by direct computation and the proof is omitted here.

The following lemma, which gives a powerful criterion of the polynomials of the form $x^l g(x^{\frac{p^n-1}{s}})$ to be permutation polynomials, will be used to prove our main results.

Lemma 2 *[17, 20] Let* p *be a prime. Let* l, n *and* s *be positive integers such that* $s \mid (p^n - 1)$*. Let* $g(x) \in \mathbb{F}_{p^n}[x]$*. Then* $f(x) = x^l g(x^{\frac{p^n-1}{s}})$ *is a permutation polynomial over* $\mathbb{F}_{p^n}$ *if and only if*

1) $\gcd(l, \frac{p^n-1}{s}) = 1$*, and*

2) $x^l g(x)^{\frac{p^n-1}{s}}$ *permutes* μ_s*,*

where μ_s *is the set of* s*-th roots of unity in* $\mathbb{F}_{p^n}$*.*

3 New classes of permutation trinomials with Niho exponents

In this section, we present several classes of fractional polynomials that permute the set of the $(q+1)$-th roots of unity in $\mathbb{F}_{q^2}$, then by Lemma 2, we construct several classes of permutation trinomials of the form (1).

For simplicity, throughout this section, we denote $\Omega_+ = \{x^2 \mid x \in \mu_{q+1}\}$ and $\Omega_- = \{-x^2 \mid x \in \mu_{q+1}\}$.

Lemma 3 *Let $q = 5^k$, where k is a positive integer. Let $g_1(x) = -x^{\frac{q+1}{2}}(\frac{x^s-2}{x^s+2})^2$, where $s = \frac{q+3}{4}$. Then $g_1(x)$ permutes μ_{q+1}.*

Proof: Note that $x^{\frac{q+1}{2}} = \pm 1$. In the following we show that if $x^{\frac{q+1}{2}} = 1$, i.e., $x \in \Omega_+$, then $-x^{\frac{q+1}{2}}(\frac{x^s-2}{x^s+2})^2 = -(\frac{x^s-2}{x^s+2})^2$ permutes Ω_+. Similarly, if $x^{\frac{q+1}{2}} = -1$, then $-x^{\frac{q+1}{2}}(\frac{x^s-2}{x^s+2})^2 = (\frac{x^s-2}{x^s+2})^2$ permutes Ω_-. Then the conclusion follows from $\Omega_+ \bigcup \Omega_- = \mu_{q+1}$ and $\Omega_+ \bigcap \Omega_- = \emptyset$.

By $x^{\frac{q+1}{2}} = 1$, i.e., $x \in \Omega_+$, we have

$$(-(\frac{x^s-2}{x^s+2})^2)^{\frac{q+1}{2}} = -(\frac{x^s-2}{x^s+2})^{q+1} = -(\frac{x^{-s}-2}{x^{-s}+2})(\frac{x^s-2}{x^s+2}) = -(\frac{2+x^s}{2-x^s})(\frac{x^s-2}{x^s+2}) = 1,$$

which shows that if $x \in \Omega_+$, then $-(\frac{x^s-2}{x^s+2})^2 \in \Omega_+$. Suppose that there exist two different elements $x_1 = y_1^2, x_2 = y_2^2 \in \Omega_+$ such that $-(\frac{x_1^s-2}{x_1^s+2})^2 = -(\frac{x_2^s-2}{x_2^s+2})^2$. Then $\frac{x_1^s-2}{x_1^s+2} = \pm\frac{x_2^s-2}{x_2^s+2}$.

1) If $\frac{x_1^s-2}{x_1^s+2} = \frac{x_2^s-2}{x_2^s+2}$, then $1 + \frac{1}{x_1^s+2} = 1 + \frac{1}{x_2^s+2}$, i.e., $x_1^s = x_2^s$. Since $x_1^s = y_1^{2s} = y_1^{\frac{q+1}{2}+1}$ and $x_2^s = y_2^{2s} = y_2^{\frac{q+1}{2}+1}$, we have $y_1 = \pm y_2$ due to $y_1^{\frac{q+1}{2}} = \pm 1$ and $y_2^{\frac{q+1}{2}+1} = \pm 1$. Therefore, $x_1 = x_2$.

2) If $\frac{x_1^s-2}{x_1^s+2} = -\frac{x_2^s-2}{x_2^s+2}$, then $2 + \frac{1}{2+x_1^s} + \frac{1}{2+x_2^s} = 0$, i.e.,

$$2(2+x_1^s)(2+x_2^s) + 2 + x_1^s + 2 + x_2^s = 0. \tag{2}$$

By (2), we get $x_1^s x_2^s = -1$. Note that $x_1^s = y_1^{2s} = y_1^{\frac{q+1}{2}+1} = \pm y_1$ and $x_2^s = y_2^{2s} = y_2^{\frac{q+1}{2}+1} = \pm y_2$, we obtain $y_1 y_2 = \pm 1$, thus $x_1 x_2 = y_1^2 y_2^2 = 1$, a contradiction with $(x_1 x_2)^s = -1$.

Based on the above analysis, $-(\frac{x^s-2}{x^s+2})^2$ permutes Ω_+. □

Theorem 1 *Let $q = 5^k$, $s = \frac{q+3}{4}$, and $t = 2s = \frac{q+3}{2}$. Then $x + x^{s(q-1)+1} - x^{t(q-1)+1}$ permutes $\mathbb{F}_{q^2}$.*

Proof: According to Lemma 2, it is sufficient to show that $x(1 + x^s - x^t)^{q-1}$ permutes μ_{q+1}. First we prove that $1 + x^s - x^t \neq 0$ for any $x \in \mu_{q+1}$. Otherwise, $1 + x^s - x^t = -(x^s + 2)^2 = 0$, i.e., $x^s = -2$. However, $(x^s)^{q+1} = (-2)^{q+1} = -1$, a contradiction with $x \in \mu_{q+1}$.

Now suppose that $1+x^s-x^t \neq 0$ for any $x \in \mu_{q+1}$. Thus,

$$\begin{aligned} x(1+x^s-x^t)^{q-1} &= x\frac{(1+x^s-x^{2s})^q}{1+x^s-x^{2s}} = \frac{x(1+x^{-s}-x^{-2s})}{1+x^s-x^{2s}} \\ &= -x^{1-2s}(\frac{x^s-2}{x^s+2})^2 = -x^{-\frac{q+1}{2}}(\frac{x^s-2}{x^s+2})^2. \end{aligned}$$

Then the conclusion follows from Lemma 3 and $x^{-\frac{q+1}{2}} = x^{\frac{q+1}{2}}$. □

Corollary 1 *Let $q=5^k$. Then $f(x)=x^q+x^{(\frac{3}{4}(q-1)+2)(q-1)+1}-x^{\frac{q+1}{2}(q-1)+1}$ is a permutation trinomial over $\mathbb{F}_{q^2}$.*

Proof: According to Lemma 2, it is sufficient to prove that $g_2(x)=x(x+x^{\frac{3}{4}(q-1)+2}-x^{\frac{q+1}{2}})^{q-1}$ permutes μ_{q+1}. Note that $x^{-\frac{q+1}{2}} = x^{\frac{q+1}{2}}$, we have

$$\begin{aligned} g_2(x) &= x(x+x^{\frac{3}{4}(q-1)+2}-x^{\frac{q+1}{2}})^{q-1} = x\frac{x^{-1}+x^{-\frac{3}{4}(q-1)-2}-x^{-\frac{q+1}{2}}}{x+x^{\frac{3}{4}(q-1)+2}-x^{\frac{q+1}{2}}} \\ &= \frac{1+x^{q+1}\cdot x^{-\frac{3}{4}(q-1)-1}-x^{q+1}\cdot x^{-\frac{q-1}{2}}}{x+x^{\frac{3}{4}(q-1)+2}-x^{\frac{q+1}{2}}} = \frac{1+x^{\frac{1}{4}(q-1)+1}-x^{\frac{q+1}{2}+1}}{x+x^{\frac{3}{4}(q-1)+2}-x^{\frac{q+1}{2}}} \\ &= \frac{(1+x^{\frac{1}{4}(q-1)+1}-x^{\frac{q+1}{2}+1})\cdot x^{\frac{q+1}{2}}}{(x+x^{\frac{3}{4}(q-1)+2}-x^{\frac{q+1}{2}})\cdot x^{\frac{q+1}{2}}} = -\frac{x^{\frac{q+1}{2}}(x^{\frac{q+3}{4}}+2)^2}{(x^{\frac{q+3}{4}}-2)^2} \\ &= -\frac{x^{-\frac{q+1}{2}}(x^{\frac{q+3}{4}}+2)^2}{(x^{\frac{q+3}{4}}-2)^2} = \frac{1}{g_1(x)}. \end{aligned}$$

Then the conclusion follows from Lemma 3. □

Lemma 4 *Let $q=5^k$, where k is odd. Let $f(x)=-x(\frac{x-2}{x+2})^2$ and $g(x)=-x(\frac{x+2}{x-2})^2$. Then $f(x)$ permutes Ω_+, $g(x)$ permutes Ω_-.*

Proof: We will show that $f(x)$ permutes Ω_+. Then $g(x)$ permutes Ω_- can be proved in a similar way.

Obviously, if $x \in \Omega_+$, then $(-x(\frac{x-2}{x+2})^2)^{\frac{q+1}{2}} = -(\frac{x-2}{x+2})^{q+1} = -(\frac{x-2}{x+2})^q \cdot \frac{x-2}{x+2} = -\frac{x^{-1}-2}{x^{-1}+2}\cdot\frac{x-2}{x+2} = 1$, which shows that $-x(\frac{x-2}{x+2})^2 \in \Omega_+$, i.e., $\{-x(\frac{x-2}{x+2})^2 \mid x \in \Omega_+\} \subseteq \Omega_+$. Suppose that there exist two different elements $x_1=y_1^2, x_2=y_2^2 \in \Omega_+$ such that $-x_1(\frac{x_1-2}{x_1+2})^2 = -x_2(\frac{x_2-2}{x_2+2})^2$. Then $y_1(\frac{y_1^2-2}{y_1^2+2}) = \pm y_2(\frac{y_2^2-2}{y_2^2+2})$. Let $y_2 = uy_1$ for some $u \in \mu_{q+1}$.

1) If $y_1(\frac{y_1^2-2}{y_1^2+2}) = y_2(\frac{y_2^2-2}{y_2^2+2})$, then $(-y_1^5-2y_1^3)u^3+(y_1^5-2y_1^3)u^2+(2y_1^3-y_1)u+2y_1^3+y_1=0$, i.e.,

$$-y_1(u-1)((y_1^4+2y_1^2)u^2-y_1^2u+2y_1^2+1)=0.$$

Since $y_1 \neq 0$, then $u = 1$ or $(y_1^4 + 2y_1^2)u^2 - y_1^2 u + 2y_1^2 + 1 = 0$. Suppose that $(y_1^4 + 2y_1^2)u^2 - y_1^2 u + 2y_1^2 + 1 = 0$. Notice that the discriminant of $(y_1^4 + 2y_1^2)u^2 - y_1^2 u + 2y_1^2 + 1$ is $\Delta = 2y_1^2(y_1^2 - 1)^2$, then

$$u = \frac{y_1^2 \pm \sqrt{2} y_1 (y_1^2 - 1)}{2y_1^2(y_1^2 + 2)}.$$

Since $q = 5^k$ and k is odd, we have $(\sqrt{2})^{q-1} = 2^{\frac{q-1}{2}} = (-1)^{\frac{5^k-1}{5-1}} = -1$. Thus $\sqrt{2}^q = -\sqrt{2}$. By $u^{q+1} = (\frac{y_1^2 \pm \sqrt{2} y_1 (y_1^2 - 1)}{2y_1^2(y_1^2+2)})^{q+1} = 1$, we have

$$\begin{aligned} (\frac{y_1^2 \pm \sqrt{2} y_1 (y_1^2 - 1)}{2y_1^2(y_1^2 + 2)})^{q+1} &= (\frac{y_1^2 \pm \sqrt{2} y_1 (y_1^2 - 1)}{2y_1^2(y_1^2 + 2)})^{q} (\frac{y_1^2 \pm \sqrt{2} y_1 (y_1^2 - 1)}{2y_1^2(y_1^2 + 2)}) \\ &= (\frac{y_1^{-2} \mp \sqrt{2} y_1^{-1} (y_1^{-2} - 1)}{2y_1^{-2}(y_1^{-2} + 2)}) (\frac{y_1^2 \pm \sqrt{2} y_1 (y_1^2 - 1)}{2y_1^2(y_1^2 + 2)}) = 1, \end{aligned}$$

then it follows that

$$(y_1^2 - 1)^2 = \pm 2\sqrt{2} y_1 (y_1^2 - 1). \tag{3}$$

If $y_1^2 = 1$, then $u = \frac{y_1^2 \pm \sqrt{2} y_1 (y_1^2 - 1)}{2y_1^2(y_1^2+2)} = 1$. Otherwise, take the 2-th power on both sides of (3), we have $y_1^4 = -1$, i.e., $x_1 = y_1^2 = \pm 2$, which is a contradiction since $\pm 2 \notin \mu_{q+1}$. As a result, $u = 1$, i.e., $x_1 = x_2$.

2) If $y_1(\frac{y_1^2 - 2}{y_1^2 + 2}) = -y_2(\frac{y_2^2 - 2}{y_2^2 + 2})$, then $(y_1^5 + 2y_1^3)u^3 + (y_1^5 + 3y_1^3)u^2 + (y_1 + 3y_1^3)u + 2y_1^3 + y_1 = 0$, i.e.,

$$y_1(u + 1)((y_1^4 + 2y_1^2)u^2 + y_1^2 u + 2y_1^2 + 1) = 0.$$

Since $y_1 \neq 0$, then $u = -1$ or $(y_1^4 + 2y_1^2)u^2 + y_1^2 u + 2y_1^2 + 1 = 0$. Suppose that $(y_1^4 + 2y_1^2)u^2 + y_1^2 u + 2y_1^2 + 1 = 0$. Notice that the discriminant of $(y_1^4 + 2y_1^2)u^2 + y_1^2 u + 2y_1^2 + 1$ is $\Delta = 2y_1^2(y_1^2 - 1)^2$, then

$$u = \frac{-y_1^2 \pm \sqrt{2} y_1 (y_1^2 - 1)}{2y_1^2(y_1^2 + 2)}.$$

Similar as in Case 1), we have

$$(y_1^2 - 1)^2 = \mp 2\sqrt{2} y_1 (y_1^2 - 1).$$

If $y_1^2 = 1$, then $u = -1$. Otherwise, take the 2-th power on both sides of $y_1^2 - 1 = \mp 2\sqrt{2} y_1$, we have $y_1^4 = -1$, i.e., $x_1 = y_1^2 = \pm 2$, which is a contradiction since $\pm 2 \notin \mu_{q+1}$. As a result, $u = -1$, i.e., $y_1 = -y_2$ and $x_1 = x_2$.

Based on the above analysis, we conclude that $-x(\frac{x-2}{x+2})^2$ permutes Ω_+. $\square$

Theorem 2 *Let $q = 5^k$, where k is odd. Let $s = \frac{q-1}{2}$ and $t = \frac{q+3}{2}$. Then $x + x^{s(q-1)+1} - x^{t(q-1)+1}$ permutes $\mathbb{F}_{q^2}$.*

Proof: According to Lemma 2, it is sufficient to prove that $g_3(x) = x(1 + x^{\frac{q-1}{2}} - x^{\frac{q+3}{2}})^{q-1}$ permutes μ_{q+1}. If $1 + x^{\frac{q-1}{2}} - x^{\frac{q+3}{2}} = 0$ for some $x \in \mu_{q+1}$, then $x + x^{\frac{q+1}{2}} - x \cdot x^{\frac{q+3}{2}} = 0$. By $x^{\frac{q+1}{2}} = \pm 1$, we have $x + x^{\frac{q+1}{2}} - x \cdot x^{\frac{q+3}{2}} = x \pm 1 \mp x^2 = 0$, which means that $x = \mp 2$, a contradiction with $x \in \mu_{q+1}$ since $\pm 2 \notin \mu_{q+1}$. Thus, $1 + x^{\frac{q-1}{2}} - x^{\frac{q+3}{2}} \neq 0$ for any $x \in \mu_{q+1}$. Then we have

$$\begin{aligned} g_3(x) = x(1 + x^{\frac{q-1}{2}} - x^{\frac{q+3}{2}})^{q-1} &= x\frac{(1 + x^{\frac{q-1}{2}} - x^{\frac{q+3}{2}})^q}{1 + x^{\frac{q-1}{2}} - x^{\frac{q+3}{2}}} \\ &= \frac{x(1 + x^{-\frac{q-1}{2}} - x^{-\frac{q+3}{2}})}{1 + x^{\frac{q-1}{2}} - x^{\frac{q+3}{2}}} \\ &= \frac{x^2(x^{-1} + x^{-\frac{q+1}{2}} - x^{-2}x^{-\frac{q+1}{2}})}{1 + x^{-1}x^{\frac{q+1}{2}} - x \cdot x^{\frac{q+1}{2}}}. \end{aligned}$$

If $x \in \Omega_+$, i.e., $x^{\frac{q+1}{2}} = 1$, then $g_3(x) = x(1 + x^{\frac{q-1}{2}} - x^{\frac{q+3}{2}})^{q-1} = \frac{x^2(x^{-1}+x^{-\frac{q+1}{2}}-x^{-2}x^{-\frac{q+1}{2}})}{1+x^{-1}x^{\frac{q+1}{2}}-x\cdot x^{\frac{q+1}{2}}} = \frac{x^2(x^{-1}+1-x^{-2})}{1+x^{-1}-x} = -x(\frac{x-2}{x+2})^2$. By Lemma 4, $g_3(x)$ permutes Ω_+.

If $x \in \Omega_-$, i.e., $x^{\frac{q+1}{2}} = -1$, then $g_3(x) = x(1 + x^{\frac{q-1}{2}} - x^{\frac{q+3}{2}})^{q-1} = \frac{x^2(x^{-1}+x^{-\frac{q+1}{2}}-x^{-2}x^{-\frac{q+1}{2}})}{1+x^{-1}x^{\frac{q+1}{2}}-x\cdot x^{\frac{q+1}{2}}} = \frac{x^2(x^{-1}-1+x^{-2})}{1-x^{-1}+x} = -x(\frac{x+2}{x-2})^2$. Again by Lemma 4, $g_3(x)$ permutes Ω_-. Then by $\Omega_+ \bigcup \Omega_- = \mu_{q+1}$ and $\Omega_+ \bigcap \Omega_- = \emptyset$, we have $g_3(x) = x(1 + x^{\frac{q-1}{2}} - x^{\frac{q+3}{2}})^{q-1}$ permutes μ_{q+1}. This completes the proof. $\square$

Corollary 2 *Let $q = 5^k$, where k is an odd integer. Then $f(x) = x^q + x^{\frac{q+5}{2}(q-1)+1} - x^{\frac{q+1}{2}(q-1)+1}$ is a permutation trinomial over $\mathbb{F}_{q^2}$.*

Proof: According to Lemma 2, it is sufficient to prove that $g_4(x) = x(x + x^{\frac{q+5}{2}} - x^{\frac{q+1}{2}})^{q-1}$ permutes μ_{q+1}. Note that

$$\begin{aligned} g_4(x) &= x(x + x^{\frac{q+5}{2}} - x^{\frac{q+1}{2}})^{q-1} = x\frac{x^{-1} + x^{-\frac{q+1}{2}-2} - x^{-\frac{q+1}{2}}}{x + x^{\frac{q+1}{2}+2} - x^{\frac{q+1}{2}}} \\ &= x\frac{x^{-1} + x^{\frac{q+1}{2}-2} - x^{\frac{q+1}{2}}}{x + x^{-\frac{q+1}{2}+2} - x^{-\frac{q+1}{2}}} = x\frac{x^{-1} + x^{\frac{q+1}{2}-2} - x^{\frac{q+1}{2}}}{x(1 + x^{-\frac{q+1}{2}+1} - x^{-\frac{q+3}{2}})} \\ &= \frac{1 + x^{\frac{q+1}{2}-1} - x^{\frac{q+3}{2}}}{x(1 + x^{-\frac{q+1}{2}+1} - x^{-\frac{q+3}{2}})} = \frac{1}{g_3(x)}, \end{aligned}$$

where the third equal sign holds due to $x^{-\frac{q+1}{2}} = x^{\frac{q+1}{2}}$. By Theorem 2, $g_3(x)$ permutes μ_{q+1} if k is odd. This completes the proof. □

Theorem 3 *Let $q = 5^k$, where k is odd. Then the following trinomials permute $\mathbb{F}_{q^2}$:*

1) $x - x^{\frac{(q+3)(q-1)}{2}+1} + x^{q(q-1)+1}$, and

2) $x^q - x^{\frac{q+1}{2}(q-1)+1} + x^{2(q-1)+1}$.

Proof: 1) According to Lemma 2, we only need to show that $g_5(x) = x(1-x^{\frac{q+3}{2}}+x^{-1})^{q-1}$ permutes μ_{q+1}.

Suppose that $x \in \Omega_+$, i.e., $x^{\frac{q+1}{2}} = 1$, then $g_5(x) = x(1 - x^{\frac{q+3}{2}} + x^{-1})^{q-1} = x(1 - x + x^{-1})^{q-1} = x \cdot \frac{1-x^{-1}+x}{1-x+x^{-1}}$ due to $1 - x + x^{-1} \neq 0$ for $x \in \Omega_+$. Otherwise, we have $x(1 - x + x^{-1}) = -(x^2 - x - 1) = -(x + 2)^2 = 0$, i.e., $x = -2$, a contradiction with $-2 \notin \Omega_+$. Therefore $g_5(x) = x \cdot \frac{1-x^{-1}+x}{1-x+x^{-1}} = -x \cdot \frac{x^2+x-1}{x^2-x-1} = -x(\frac{x-2}{x+2})^2$. By Lemma 4, $g_5(x)$ permutes Ω_+.

Suppose that $x \in \Omega_-$, i.e., $x^{\frac{q+1}{2}} = -1$, then $g_5(x) = x(1 - x^{\frac{q+3}{2}} + x^{-1})^{q-1} = x(1 + x + x^{-1})^{q-1} = x \cdot \frac{1+x^{-1}+x}{1+x+x^{-1}} = x$ due to $1+x+x^{-1} \neq 0$ for $x \in \Omega_-$. Otherwise, we have $x(1+x+x^{-1}) = x^2+x+1 = 0$, then $(x-1)(x^2+x+1) = x^3 - 1 = 0$ for some $x = -y^2 \in \Omega_-$, where $y \in \mu_{q+1}$. Thus, $y^6 = -1$ and $y^{12} = 1$, together with $y^{q+1} = 1$ and $\gcd(q+1, 12) \mid 6$, we have $y^6 = 1$, a contradiction. Therefore, $g_5(x) = x$ permutes Ω_-.

Based on the above analysis, we conclude that $g_5(x) = x(1 - x^{\frac{q+3}{2}} + x^{-1})^{q-1} = x \cdot \frac{1-x^{-\frac{q+3}{2}}+x}{1-x^{\frac{q+3}{2}}+x^{-1}}$ permutes μ_{q+1}.

2) By Lemma 2, we only need to show that $g_6(x) = x(x - x^{\frac{q+1}{2}} + x^2)^{q-1}$ permutes μ_{q+1}. Since $x^{\frac{q+1}{2}} = x^{-\frac{q+1}{2}}$, we have

$$\begin{aligned} g_6(x) &= x \cdot \frac{x^{-1} - x^{-\frac{q+1}{2}} + x^{-2}}{x - x^{\frac{q+1}{2}} + x^2} = x \cdot \frac{x^{-1} - x^{\frac{q+1}{2}} + x^{-2}}{x - x^{-\frac{q+1}{2}} + x^2} \\ &= \frac{1 - x^{\frac{q+3}{2}} + x^{-1}}{x(1 - x^{-\frac{q+3}{2}} + x)} = \frac{1}{g_5(x)} \end{aligned}$$

Therefore, $g_6(x)$ permutes μ_{q+1}. □

Similar as the proof of Theorem 3, we have

Theorem 4 *Let $q = 5^k$, where k is odd. Then the following two trinomials permute $\mathbb{F}_{q^2}$:*

1) $x - x^{s(q-1)+1} + x^{t(q-1)+1}$, where $s = \frac{q+3}{2}$ and $t = \frac{q+5}{2}$, and

2) $x^q - x^{s(q-1)+1} + x^{t(q-1)+1}$, *where* $s = \frac{q+1}{2}$ *and* $t = \frac{q-1}{2}$.

Lemma 5 *Let* $q = 5^k$, *where* k *is even. Let* $f(x) = -x(\frac{x-2}{x+2})^2$ *and* $g(x) = -x(\frac{x+2}{x-2})^2$. *Then* $f(x)$ *permutes* Ω_- *and* $g(x)$ *permutes* Ω_+.

Proof: In the following we will prove that $g(x)$ permutes Ω_+. Then $f(x)$ permutes Ω_- can be proved similarly. Obviously, if $x \in \Omega_+$, then $(-x(\frac{x+2}{x-2})^2)^{\frac{q+1}{2}} = -(\frac{x+2}{x-2})^{q+1} = -(\frac{x+2}{x-2})^q \cdot \frac{x+2}{x-2} = -\frac{x^{-1}+2}{x^{-1}-2} \cdot \frac{x+2}{x-2} = 1$, which shows that $-x(\frac{x+2}{x-2})^2 \in \Omega_+$, i.e., $\{-x(\frac{x+2}{x-2})^2 \mid x \in \Omega_+\} \subseteq \Omega_+$.

Suppose that there exist two different elements $x_1 = y_1^2, x_2 = y_2^2 \in \Omega_+$ such that $-x_1(\frac{x_1+2}{x_1-2})^2 = -x_2(\frac{x_2+2}{x_2-2})^2$. Then $y_1(\frac{y_1^2+2}{y_1^2-2}) = \pm y_2(\frac{y_2^2+2}{y_2^2-2})$. Let $y_2 = uy_1$ for some $u \in \mu_{q+1}$.

1) If $y_1(\frac{y_1^2+2}{y_1^2-2}) = y_2(\frac{y_2^2+2}{y_2^2-2})$, then $(4y_1^5 + 2y_1^3)u^3 + (y_1^5 + 2y_1^3)u^2 + (3y_1^3 + 4y_1)u + 3y_1^3 + y_1 = 0$, i.e.,

$$-y_1(u-1)((y_1^4 - 2y_1^2)u^2 + y_1^2 u - 2y_1^2 + 1) = 0.$$

Since $y_1 \neq 0$, then $u = 1$ or $(y_1^4 - 2y_1^2)u^2 + y_1^2 u - 2y_1^2 + 1 = 0$. Suppose that $(y_1^4 - 2y_1^2)u^2 + y_1^2 u - 2y_1^2 + 1 = 0$. Notice that the discriminant of $(y_1^4 - 2y_1^2)u^2 + y_1^2 u - 2y_1^2 + 1$ is $\Delta = -2y_1^2(y_1^2+1)^2$, then

$$u = \frac{-y_1^2 \pm 2\sqrt{2}y_1(y_1^2+1)}{2y_1^2(y_1^2-2)}.$$

Since $q = 5^k$ and k is even, we have $(\sqrt{2})^{q-1} = 2^{\frac{q-1}{2}} = (-1)^{\frac{5^k-1}{5-1}} = 1$. Thus $\sqrt{2}^q = \sqrt{2}$. By $u^{q+1} = (\frac{-y_1^2 \pm 2\sqrt{2}y_1(y_1^2+1)}{2y_1^2(y_1^2-2)})^{q+1} = 1$, we have

$$\begin{aligned}(\frac{-y_1^2 \pm 2\sqrt{2}y_1(y_1^2+1)}{2y_1^2(y_1^2-2)})^{q+1} &= (\frac{-y_1^2 \pm 2\sqrt{2}y_1(y_1^2+1)}{2y_1^2(y_1^2-2)})^q(\frac{-y_1^2 \pm 2\sqrt{2}y_1(y_1^2+1)}{2y_1^2(y_1^2-2)}) \\ &= (\frac{-y_1^{-2} \pm 2\sqrt{2}y_1^{-1}(y_1^{-2}+1)}{2y_1^{-2}(y_1^{-2}-2)})(\frac{-y_1^2 \pm 2\sqrt{2}y_1(y_1^2+1)}{2y_1^2(y_1^2-2)}) = 1\end{aligned}$$

then it follows that

$$(y_1^2+1)^2 = \mp\sqrt{2}y_1(y_1^2+1). \tag{4}$$

If $y_1^2 = -1$, then $u = \frac{-y_1^2 \pm 2\sqrt{2}y_1(y_1^2+1)}{2y_1^2(y_1^2-2)} = 1$. Otherwise, take the 2-th power on both sides of (4), we have $y_1^4 = -1$, i.e., $x_1 = y_1^2 = \pm 2$, which is a contradiction since $\pm 2 \notin \mu_{q+1}$. As a result, $u = 1$, i.e., $x_1 = x_2$.

2) If $y_1(\frac{y_1^2+2}{y_1^2-2}) = -y_2(\frac{y_2^2+2}{y_2^2-2})$, then $(y_1^5+3y_1^3)u^3+(y_1^5+2y_1^3)u^2+(2y_1^3+y_1)u+3y_1^3+y_1=0$, i.e.,

$$y_1(u+1)((y_1^4-2y_1^2)u^2-y_1^2u-2y_1^2+1)=0.$$

Since $y_1 \neq 0$, then $u=-1$ or $(y_1^4-2y_1^2)u^2-y_1^2u-2y_1^2+1=0$. Suppose that $(y_1^4-2y_1^2)u^2-y_1^2u-2y_1^2+1=0$. Notice that the discriminant of $(y_1^4-2y_1^2)u^2-y_1^2u-2y_1^2+1$ is $\Delta=-2y_1^2(y_1^2+1)^2$, then

$$u=\frac{y_1^2 \pm 2\sqrt{2}y_1(y_1^2+1)}{2y_1^2(y_1^2-2)}.$$

Similar as in Case 1), we have

$$(y_1^2+1)^2 = \pm\sqrt{2}y_1(y_1^2+1).$$

If $y_1^2=-1$, then $u=\frac{y_1^2\pm 2\sqrt{2}y_1(y_1^2+1)}{2y_1^2(y_1^2-2)}=-1$. Otherwise, take the 2-th power on both sides of $(y_1^2+1)=\pm\sqrt{2}y_1$, we have $y_1^4=-1$, i.e., $x_1=y_1^2=\pm 2$, which is a contradiction since $\pm 2 \notin \mu_{q+1}$. As a result, $u=-1$, i.e., $y_1=-y_2$ and $x_1=x_2$.

Therefore, $-x(\frac{x+2}{x-2})^2$ permutes Ω_+. □

Theorem 5 *Let $q=5^k$, where k is even. Then the following two trinomials permute $\mathbb{F}_{q^2}$:*

1) $x-x^{2(q-1)+1}+x^{\frac{(q+3)(q-1)}{2}+1}$, *and*

2) $x^q-x^{q^2-q+1}+x^{\frac{q+1}{2}(q-1)+1}$.

Proof: 1) According to Lemma 2, we only need to show that $g_7(x)=x(1-x^2+x^{\frac{q+3}{2}})^{q-1}$ permutes μ_{q+1}. First we show that $1-x^2+x^{\frac{q+3}{2}}=1-x^2\pm x\neq 0$ for any $x\in\mu_{q+1}$. Otherwise, $1-x^2\pm x=-(x\pm 2)^2=0$, i.e., $x=\mp 2$, a contradiction with $\pm 2\notin\mu_{q+1}$.

Suppose that $x\in\Omega_+$, i.e., $x^{\frac{q+1}{2}}=1$, then

$$\begin{aligned} g_7(x) &= x(1-x^2+x^{\frac{q+3}{2}})^{q-1} = x(1-x^2+x)^{q-1} \\ &= x\cdot\frac{1-x^{-2}+x^{-1}}{1-x^2+x} = -x^{-1}\cdot\frac{x^2+x-1}{x^2-x-1} = -x^{-1}(\frac{x-2}{x+2})^2, \end{aligned}$$

thus $g_7(x)$ permutes Ω_+ by Lemma 5.

Suppose that $x\in\Omega_-$, i.e., $x^{\frac{q+1}{2}}=-1$, then

$$\begin{aligned} g_7(x) &= x(1-x^2+x^{\frac{q+3}{2}})^{q-1} = x(1-x^2-x)^{q-1} \\ &= x\cdot\frac{1-x^{-2}-x^{-1}}{1-x^2-x} = -x^{-1}\cdot\frac{x^2-x-1}{x^2+x-1} = -x^{-1}(\frac{x+2}{x-2})^2. \end{aligned}$$

Again by Lemma 5, $g_7(x)$ permutes Ω_-.

Based on the above analysis, we conclude that $g_7(x) = x(1 - x^2 + x^{\frac{q+3}{2}})^{q-1} = x \cdot \frac{1-x^{-2}+x^{-\frac{q+3}{2}}}{1-x^2+x^{\frac{q+3}{2}}}$ permutes μ_{q+1}.

2) By Lemma 2, we only need to show that $g_8(x) = x(x - x^{-1} + x^{\frac{q+1}{2}})^{q-1}$ permutes μ_{q+1}. Since $x^{\frac{q+1}{2}} = x^{-\frac{q+1}{2}}$, we have

$$\begin{aligned} g_8(x) &= x \cdot \frac{x^{-1} - x + x^{\frac{q+1}{2}}}{x - x^{-1} + x^{\frac{q+1}{2}}} = \frac{1 - x^2 + x^{\frac{q+3}{2}}}{x - x^{-1} + x^{\frac{q+1}{2}}} \\ &= \frac{1 - x^2 + x^{\frac{q+3}{2}}}{x(1 - x^{-2} + x^{\frac{q+1}{2}-1})} = \frac{1 - x^2 + x^{\frac{q+3}{2}}}{x(1 - x^{-2} + x^{-\frac{q+1}{2}-1})} = \frac{1}{g_7(x)} \end{aligned}$$

Therefore, $g_8(x)$ permutes μ_{q+1}. □

Theorem 6 *Let $q = 5^k$, where k is even. Let $s = 1$ and $t = \frac{q-1}{2}$. Then $x + x^{s(q-1)+1} - x^{t(q-1)+1}$ permutes $\mathbb{F}_{q^2}$.*

Proof: According to Lemma 2, it is sufficient to prove that $g_9(x) = x(1 + x^s - x^t)^{q-1}$ permutes μ_{q+1}. First we show that $1 + x^s - x^t \neq 0$ for any $x \in \mu_{q+1}$.

Suppose that $1 + x^s - x^t = 1 + x - x^{\frac{q-1}{2}} = 0$ for some $x \in \mu_{q+1}$, then $x + x^2 - x^{\frac{q+1}{2}} = 0$. Note that $x^{\frac{q+1}{2}} = \pm 1$. Thus $x + x^2 \pm 1 = 0$. If $x + x^2 - 1 = (x-2)^2 = 0$, then $x = 2$, a contradiction with $x \in \mu_{q+1}$ since $2 \notin \mu_{q+1}$. If $x + x^2 + 1 = 0$, then $x^3 = 1$. Together with $x^{q+1} = 1$ and $\gcd(3, q+1) = 1$ due to k is even, we have $x = 1$, then we have $0 = x + x^2 + 1 = 3$, a contradiction. Therefore, $1 + x - x^{\frac{q-1}{2}} \neq 0$ for any $x \in \mu_{q+1}$.

Then we have

$$\begin{aligned} g_9(x) = x(1 + x - x^{\frac{q-1}{2}})^{q-1} &= x\frac{(1 + x - x^{\frac{q-1}{2}})^q}{1 + x - x^{\frac{q-1}{2}}} \\ &= \frac{x(1 + x^{-1} - x^{-\frac{q-1}{2}})}{1 + x - x^{\frac{q-1}{2}}} \\ &= \frac{x^2(x^{-1} + x^{-2} - x^{-\frac{q+1}{2}})}{1 + x - x^{-1} \cdot x^{\frac{q+1}{2}}}. \end{aligned}$$

If $x \in \Omega_-$, i.e., $x^{\frac{q+1}{2}} = -1$, then $x(1 + x - x^{\frac{q-1}{2}})^{q-1} = \frac{x^2(x^{-1}+x^{-2}-x^{-\frac{q+1}{2}})}{1+x-x^{-1}\cdot x^{\frac{q+1}{2}}} = \frac{x^2(x^{-1}+x^{-2}+1)}{1+x+x^{-1}} = x$ permutes Ω_-.

If $x \in \Omega_+$, i.e., $x^{\frac{q+1}{2}} = 1$, then by Lemma 5, $x(1+x-x^{\frac{q-1}{2}})^{q-1} = \frac{x^2(x^{-1}+x^{-2}-x^{-\frac{q+1}{2}})}{1+x-x^{-1}\cdot x^{\frac{q+1}{2}}} = \frac{x^2(x^{-1}+x^{-2}-1)}{1+x-x^{-1}} = -x(\frac{x+2}{x-2})^2$ permutes Ω_+. Then $g_9(x) = x(1+x-x^{\frac{q-1}{2}})^{q-1}$ permutes μ_{q+1} due to $\Omega_+ \bigcup \Omega_- = \mu_{q+1}$ and $\Omega_+ \bigcap \Omega_- = \emptyset$. This completes the proof. □

Similar as the proof of Theorem 6, by using Lemma 5, we can construct the following permutation trinomials over $\mathbb{F}_{5^n}$.

Theorem 7 *Let $q = 5^k$, where k is even. Then the following trinomials permute $\mathbb{F}_{q^2}$:*

1) $x - x^q + x^{\frac{q+5}{2}\cdot(q-1)+1}$,

2) $x + x^{\frac{q+3}{2}\cdot(q-1)+1} + x^{\frac{q+5}{2}\cdot(q-1)+1}$, *and*

3) $x + x^{\frac{q+3}{2}\cdot(q-1)+1} - x^{q^2-q+1}$.

It is known that the inverse of a Niho exponent, if exists, is again a Niho exponent, and the product of two Niho exponents is still a Niho exponent [3]. Similar as Lemma 4 in [12], we have the following lemma.

Lemma 6 *Let $q = p^k$, where p is a prime and k is an integer.*

1) Suppose that $f(x) = x(1+x^i+x^j)^{q-1}$ permutes μ_{q+1}. If $\gcd(2i-1, q+1) = 1$, then $x(1+x^s+x^t)^{q-1}$ permutes μ_{q+1}, where $s = \frac{i}{2i-1}$ and $t = \frac{i-j}{2i-1}$.

2) Suppose that $f(x) = x(1+x^i-x^j)^{q-1}$ permutes μ_{q+1}. If $\gcd(2i-1, q+1) = 1$, then $x(1+x^s-x^t)^{q-1}$ permutes μ_{q+1}, where $s = \frac{i}{2i-1}$ and $t = \frac{i-j}{2i-1}$; if $\gcd(2j-1, q+1) = 1$, then $x(1-x^s-x^t)^{q-1}$ permutes μ_{q+1}, where $s = \frac{j}{2j-1}$ and $t = \frac{j-i}{2j-1}$.

3) Suppose that $f(x) = x(1-x^i-x^j)^{q-1}$ permutes μ_{q+1}. If $\gcd(2i-1, q+1) = 1$, then $x(1-x^s+x^t)^{q-1}$ permutes μ_{q+1}, where $s = \frac{i}{2i-1}$ and $t = \frac{i-j}{2i-1}$.

The integers s and t written as fractions in Lemma 6 should be interpreted as modulo p^k+1. Based on Lemma 6, one can obtain lots of permutation trinomials over $\mathbb{F}_{5^n}$ from the permutation trinomials constructed in this section.

In Table 1, we list all the permutation trinomials of the form (1) constructed in this paper. We denote by $(\pm[c_1], \pm[c_2])$ the permutation trinomial $f(x) = x \pm x^{c_1(5^k-1)+1} \pm x^{c_2(5^k-1)+1}$. For example, the pair $(+[1], -[\frac{5^k-1}{2}])$ means the trinomial $x + x^{5^k} - x^{\frac{5^k-1}{2}(5^k-1)+1}$. The "Equivalent Pairs" column in Table 1 are obtained based on Lemma 6.

To end this section, we propose two conjectures. If the conjectures are true, then we can construct two classes of permutation trinomials over $\mathbb{F}_{5^n}$.

Table 1: Pairs $(\pm[c_1], \pm[c_2])$ such that $f(x)$ defined by (1) are permutation trinomials over $\mathbb{F}_{q^2}$ $(q = 5^k)$

$(\pm[c_1], \pm[c_2])$	$g(x)$	Conditions	Equivalent Pairs	Proven in
$(+[\frac{q+3}{4}], -[\frac{q+3}{2}])$	$-x^{\frac{q+1}{2}}(\frac{x^s-2}{x^s+2})^2$ $(s = \frac{q+3}{4})$	for all k	$(-[\frac{q+3}{4}], -[\frac{q+3}{2}])$	Theorem 1
$(+[\frac{q-1}{2}], -[\frac{q+3}{2}])$	$\frac{-x(x^{\frac{q+3}{2}}-x^{\frac{q-1}{2}}+1)}{x^{\frac{q+3}{2}}-x^{\frac{q-1}{2}}-1}$	k odd	$(-[2], -[\frac{q+3}{2}])$	Theorem 2
$(+[-1], -[\frac{q+3}{2}])$	$\frac{x^2(x^{\frac{q-1}{2}}-x-1)}{x^{\frac{q+5}{2}}-x-1}$	k odd	$(-[\frac{q+3}{2}], -[\frac{q+5}{2}])$	Theorem 3
$(-[\frac{q+3}{2}], +[\frac{q+5}{2}])$	$\frac{-x(x^{\frac{q-1}{2}}-x^{\frac{q-3}{2}}-1)}{x^{\frac{q+5}{2}}-x^{\frac{q+3}{2}}+1}$	k odd	$(-[-1], -[\frac{q+3}{2}])$	Theorem 4
$(-[2], +[\frac{q+3}{2}])$	$\frac{x^{\frac{q+3}{2}}+x^2-1}{x(x^{\frac{q+3}{2}}-x^2+1)}$	k even	$(+[\frac{q+3}{2}], -[\frac{q-1}{2}])$, $(-[2\cdot\frac{q+2}{3}], -[\frac{q+2}{3}\cdot\frac{q+3}{2}])$	Theorem 5
$(+[1], -[\frac{q-1}{2}])$	$\frac{x^{\frac{q+5}{2}}-x-1}{x^{\frac{q-1}{2}}-x-1}$	k even	$(+[1], -[\frac{q+5}{2}])$, $(-[\frac{q+2}{3}\cdot\frac{q+5}{2}], -[\frac{q+2}{3}\cdot\frac{q+3}{2}])$	Theorem 6
$(-[1], +[\frac{q+5}{2}])$	$\frac{x^{\frac{q-1}{2}}+x-1}{x^{\frac{q+5}{2}}-x+1}$	k even	$(-[1], -[\frac{q-1}{2}])$, $(+[\frac{q+2}{3}\cdot\frac{q+5}{2}], -[\frac{q+2}{3}\cdot\frac{q+3}{2}])$	Theorem 7
$(+[\frac{q+3}{2}], +[\frac{q+5}{2}])$	$\frac{x^{\frac{q+1}{2}}+x^{\frac{q-1}{2}}+x}{x^{\frac{q+5}{2}}+x^{\frac{q+3}{2}}+1}$	k even	$(+[\frac{q+3}{2}], +[-1])$, $(+[\frac{q+2}{3}\cdot\frac{q+5}{2}], +[\frac{q+2}{3}])$	Theorem 7
$(+[\frac{q+3}{2}], -[-1])$	$\frac{x^2(x^{\frac{q-1}{2}}-x+1)}{x^{\frac{q+5}{2}}+x-1}$	k even	$(+[\frac{q+3}{2}], -[\frac{q+5}{2}])$, $(-[\frac{q+2}{3}\cdot\frac{q+5}{2}], -[\frac{q+2}{3}])$	Theorem 7
$(+[2], -[-2])$	$-x(\frac{x^2+2}{x^2-2})^2$	k odd	$(-[-2\cdot 5^{k-1}], -[-4\cdot 5^{k-1}])$	Proposition 1
$(-[\frac{q+5}{3}], -[\frac{2(q+2)}{3}])$	$-x(\frac{x^2-2}{x^2+2})^2$	k even	$(+[-2\cdot 5^{k-1}], -[-4\cdot 5^{k-1}])$	Proposition 2

Conjecture 1 *The polynomial $x(\frac{x^2-x+2}{x^2+x+2})^2$ is a permutation polynomial over $\mathbb{F}_{5^k}$ for odd k.*

Remark 1 *Let $\Lambda_1 = \{x^2 \mid x \in \mathbb{F}_{5^k}^*\}$ and $\Lambda_2 = \{2x^2 \mid x \in \mathbb{F}_{5^k}^*\}$, where k is odd. Note that $\mathbb{F}_{5^k}^* = \Lambda_1 \bigcup \Lambda_2$ and $\Lambda_1 \bigcap \Lambda_2 = \emptyset$. Obviously, $\{x(\frac{x^2-x+2}{x^2+x+2})^2 \mid x \in \Lambda_1\} \subseteq \Lambda_1$ and $\{x(\frac{x^2-x+2}{x^2+x+2})^2 \mid x \in \Lambda_2\} \subseteq \Lambda_2$. Thus, to prove Conjecture 1 is equivalent to prove that $\{x(\frac{x^2-x+2}{x^2+x+2})^2 \mid x \in \Lambda_1\} \supseteq \Lambda_1$.*

Proposition 1 *Let $n = 2k$, where k is an odd integer. Let $q = 5^k$. If Conjecture 1 is true, then $f(x) = x + x^{2(q-1)+1} - x^{(q-1)(q-1)+1}$ permutes $\mathbb{F}_{q^2}$.*

Proof: First it is easy to show that if $x \in \mathbb{F}_{5^n} \setminus \mathbb{F}_{5^k}$, then $f(x) = x + x^{2q-1} - x^{3-2q} \in \mathbb{F}_{5^n} \setminus \mathbb{F}_{5^k}$. Otherwise, if there exists $x_1 \in \mathbb{F}_{5^n} \setminus \mathbb{F}_{5^k}$ such that $f(x_1) \in \mathbb{F}_q$, then $f(x_1) = f(x_1)^q$, i.e.,

$$x_1 + x_1^{2q-1} - x_1^{3-2q} = x_1^q + x_1^{2-q} - x_1^{3q-2}.$$

Let $y = x_1^{q-1}$, we get $1 + y^2 - y^{-2} = y + y^{-1} - y^3$, thus,

$$y^5 + y^4 - y^3 + y^2 - y - 1 = (y-1)(y^4 + 2y^3 + y^2 + 2y + 1) = 0.$$

Suppose that $y^4+2y^3+y^2+2y+1 = 0$, then the order of y is 13, thus $13 \mid (q+1)$, which is a contradiction with $\gcd(13, q+1) = 1$ due to k is odd. Therefore, we have $y = 1$, i.e., $x_1 \in \mathbb{F}_q$, a contradiction.

In the following we show that $f(x) = c$ has at most one solution for any $c \in \mathbb{F}_{q^2}$.

If $c \in \mathbb{F}_q$, then $x \in \mathbb{F}_q$, thus we have $f(x) = x = c$. Therefore, $x = c$ is the only solution of $f(x) = c$.

If $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, then $x \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, and it follows that $\mathrm{N}(x) \neq 0$. We compute

$$\begin{aligned}
\mathrm{Tr}(f(x)) &= \mathrm{Tr}(x + x^{2q-1} - x^{-2q+3}) \\
&= \mathrm{Tr}(x) + (x^{2q-1} + x^{2-q}) - (x^{-2q+3} + x^{-2+3q}) \\
&= \mathrm{Tr}(x) + \frac{x^3 + x^{3q}}{x^{q+1}} - \frac{x^5 + x^{5q}}{x^{2+2q}} \\
&= \mathrm{Tr}(x) + \frac{\mathrm{Tr}(x^3)}{\mathrm{N}(x)} - \frac{\mathrm{Tr}^5(x)}{\mathrm{N}^2(x)},
\end{aligned}$$

and

$$\begin{aligned}
\mathrm{N}(f(x)) &= \mathrm{N}(x + x^{2q-1} - x^{-2q+3}) \\
&= (x + x^{2q-1} - x^{-2q+3})(x^q + x^{2-q} - x^{-2+3q}) \\
&= 3x^{1+q} - (x^{5q-3} + x^{5-3q}) \\
&= 3\mathrm{N}(x) - \frac{x^{8q} + x^8}{x^{3q+3}} \\
&= 3\mathrm{N}(x) - \frac{\mathrm{Tr}(x^8)}{\mathrm{N}^3(x)}
\end{aligned}$$

Denote $a = \mathrm{Tr}(x)$, $b = \mathrm{N}(x)$, $\alpha = \mathrm{Tr}(f(x))$, and $\beta = \mathrm{N}(f(x))$. According to Lemma 1, we have

$$\begin{aligned}
\alpha = \mathrm{Tr}(f(x)) &= \mathrm{Tr}(x) + \frac{\mathrm{Tr}(x^3)}{\mathrm{N}(x)} - \frac{\mathrm{Tr}^5(x)}{\mathrm{N}^2(x)} \\
&= a + \frac{a^3 + 2ba}{b} - \frac{a^5}{b^2}, \\
&= a(3 + \frac{a^2}{b} - \frac{a^4}{b^2}), \qquad (5)
\end{aligned}$$

and

$$\begin{aligned}\beta = \mathrm{N}((x)) &= 3\mathrm{N}(x) - \frac{\mathrm{Tr}(x^8)}{\mathrm{N}^3(x)} \\ &= 3b - \frac{a^8 + 2ba^6 - b^3a^2 + 2b^4}{b^3} \\ &= b(1 - \frac{a^8}{b^4} - 2 \cdot \frac{a^6}{b^3} + \frac{a^2}{b}) \qquad (6)\end{aligned}$$

Denote $\gamma = \frac{\alpha^2}{\beta}$, then by (5) and (6), we have $\gamma = -r(\frac{r^2-r+2}{r^2+r+2})^2$, where $r = \frac{a^2}{b}$. If Conjecture 1 is true, we know that r is uniquely determined by γ, thus r is uniquely determined by c. From (5) and (6), it follows that $a = \mathrm{Tr}(x)$ and $b = \mathrm{N}(x)$ are uniquely determined by c. This completes the proof since (x, x^q) is uniquely determined by $a = \mathrm{Tr}(x)$ and $b = \mathrm{N}(x)$, and $f(x) \neq f(x)^q = f(x^q)$. □

Remark 2 *By Lemma 2, for odd k, $f(x) = x + x^{2(q-1)+1} - x^{(q-1)(q-1)+1}$ permutes $\mathbb{F}_{q^2}$ if and only if $g(x) = x(1 + x^2 - x^{-2})^{q-1} = x \cdot \frac{1+x^{-2}-x^2}{1+x^2-x^{-2}} = -x(\frac{x^2+2}{x^2-2})^2$ permutes μ_{q+1}. Thus, if Conjecture 1 is true, then $-x(\frac{x^2+2}{x^2-2})^2$ permutes μ_{q+1}.*

Conjecture 2 *Let $q = 5^k$, where k is an even integer. Then $-x(\frac{x^2-2}{x^2+2})^2$ permutes μ_{q+1}.*

Proposition 2 *Let $n = 2k$, where k is an even integer. Let $q = 5^k$. If Conjecture 2 is true, then $f(x) = x - x^{s(q-1)+1} - x^{t(q-1)+1}$ permutes $\mathbb{F}_{q^2}$, where $s = \frac{q+2}{3} + 1$ and $t = 2 \cdot \frac{q+2}{3}$.*

Proof: According to Lemma 2, we only need to prove that $g_{10}(x) = x(1 - x^s - x^t)^{q-1}$ permutes μ_{q+1}, which is equivalent to showing $g_{10}(x^3) = x^3(1 - x^{3s} - x^{3t})^{q-1} = x^3(1 - x^{q+2+3} - x^{2(q+2)})^{q-1} = x^3(1 - x^4 - x^2)^{q-1}$ permutes μ_{q+1} due to $\gcd(3, q+1) = 1$. Notice that $g_{10}(x^3) = x^3(1 - x^4 - x^2)^{q-1} = x^3 \cdot \frac{1-x^{-2}-x^{-4}}{1-x^2-x^4} = -x^{-1}\frac{x^4-x^2-1}{x^4+x^2-1} = -x^{-1} \cdot (\frac{x^2+2}{x^2-2})^2$. Therefore, if Conjecture 2 is true, $f(x) = x - x^{s(q-1)+1} - x^{t(q-1)+1}$ permutes $\mathbb{F}_{q^2}$. □

4 Conclusion

In this paper, we find some new fractional polynomials that permute the set of the $(q+1)$-th roots of unity in $\mathbb{F}_{q^2}$, where $q = 5^k$. Based on these fractional polynomials, we construct several classes permutation trinomials over $\mathbb{F}_{5^{2k}}$. Two conjectures are proposed, and two new classes of permutation trinomials over $\mathbb{F}_{5^{2k}}$ will be obtained if the two conjectures are true. At the end of this paper, we propose two problems about the permutation trinomials with the form (1), it will be nicer if they can be solved.

Problem 1: Let $s+t=0$ or $s+t=\frac{5^k+1}{2}$. Under what conditions for s is the polynomial $x+x^{s(5^k-1)+1}-x^{t(5^k-1)+1}$ a permutation trinomial over $\mathbb{F}_{5^{2k}}$?

Problem 2: Let $s+t=\frac{5^k+1}{2}$. Under what conditions for s is the polynomial $x+x^{s(5^k-1)+1}+x^{t(5^k-1)+1}$ a permutation trinomial over $\mathbb{F}_{5^{2k}}$?

References

[1] C. Ding, L. Qu, Q. Wang, J. Yuan, P. Yuan, Permutation trinomials over finite fields with even characteristic, SIAM J. Dis. Math. 29 (2015) 79-92.

[2] C. Ding, J. Yuan, A family of skew Hadamard difference sets, J. Comb. Theory, Ser. A 113 (2006) 1526-1535.

[3] H. Dobbertin, Almost perfect nonlinear power functions on $GF(2^n)$: the Welch case, IEEE Trans. Inf. Theory 45 (4) (1999) 1271-1275.

[4] H. Dobbertin, P. Felke, T. Helleseth, P. Rosendahl, Niho type cross-correlation functions via Dickson polynomials and Kloosterman sums, IEEE Trans. Inf. Theory 52 (2) (2006) 613-627.

[5] R. Gupta, R.K. Sharma, Some new classes of permutation trinomials over finite fields with even characteristic, Finite Fields Appl. 41 (2016) 89-96.

[6] X. Hou, Permutation polynomials over finite fields-A survey of recent advances, Finite Fields Appl. 32 (2015) 82-119.

[7] X. Hou, Determination of a type of permutaiton trinomials over finite fields, Acta Arith. 162 (2014) 253-278.

[8] X. Hou, Determination of a type of permutation trinomials over finite fields, II, Finite Fields Appl. 35 (2015) 16-35.

[9] Y. Laigle-Chapuy, Permutation polynomials and applications to coding theory, Finite Fields Appl. 13 (1) (2007) 58-70.

[10] K. Li, L. Qu, X. Chen, New classes of permutation binomials and permutation trinomials over finite fields, Finite Fields Appl. 43 (2017) 69-85.

[11] K. Li, L. Qu, C. Li, S. Fu, New permutation trinomials constructed from fractional polynomials, available online: https://arxiv.org/pdf/1605.06216v1.pdf

[12] N. Li, T. Helleseth, Several classes of permutation trinomials from Niho exponents, Cryptogr. Commun. DOI 10.1007/s12095-016-0210-9.

[13] N. Li, T. Helleseth, New permutation trinomials from Niho exponents over finite fields with even characteristic, available online: http://arxiv.org/pdf/1606.03768v1.pdf

[14] N. Li, On two conjectures about permutation trinomials over $\mathbb{F}_{3^{2k}}$, available online: http://arxiv.org/pdf/1610.04441v1.pdf

[15] R. Lidl, H. Niederreiter, Finite Fields, 2nd ed. Cambridge Univ. Press, Cambridge, 1997.

[16] Y. Niho, Multivalued cross-correlation functions between two maximal linear recursive sequences, PhD dissertation, Univ. Southern Calif. Los Angeles, 1972.

[17] Y.H. Park, J.B. Lee, Permutation polynomials and group permutation polynomials, Bull. Austral. Math. Soc. 63 (2001) 67-74.

[18] L. Qu, Y. Tan, C. H. Tan, C. Li, Constructing differentially 4-uniform permutations over $\mathbb{F}_{2^{2k}}$ via the switching method, IEEE Trans. Inf. Theory 59 (7) (2013) 4675-4686.

[19] Z. Tu, X. Zeng, L. Hu, Several classes of complete permutation polynomials, Finite Fields Appl. 25 (2014) 182-193.

[20] M.E. Zieve, On some permutation polynomials over F_q of the form $x^r h(x^{(q-1)/d})$, Proc. Amer. Math. Soc. 137 (2009), 2209-2216.

[21] M.E. Zieve, Permutation polynomials on $\mathbb{F}_q$ induced from Rédei function bijections on subgroups of $\mathbb{F}_q^*$, available online: https://arxiv.org/pdf/1310.0776v2.pdf

[22] Z. Zha, L. Hu, S. Fan, Further results on permutation trinomials over finite fields with even characteristic, Finite Fields Appl. 45 (2017) 43-52.